

# Glossary of Medical Terminology for Immunisation and Vaccine development

Supporting communication for the COVID-19 vaccination program.

### English

Produced by Health and Social Policy Branch NSW Ministry of Health, NSW Multicultural Health Communication Service, NSW Refugee Health Service and School of Population Health, University of New South Wales





#### Supporting communication for the COVID-19 vaccination program

This glossary was developed to help community organisations, translators and interpreters, bilingual workers, and community leaders to better understand and communicate about vaccine development and implementation.

#### Disclaimer

This glossary aims to provide plain language meanings to complex medical terminology about immunisation and vaccine development. The information is to be used as a reference tool only. If you would like to provide feedback or add a new word or term to the list, please contact Associate Professor Holly Seale on h.seale@unsw.edu.au or +61 (02) 9385 3129.

#### Acknowledgments

This resource was developed by:

- Associate Professor Holly Seale, School of Population Health, University of New South Wales
- Lisa Woodland, Director, NSW Multicultural Health Communication Service
- Dr Kylie Quinn, School of Health and Biomedical Sciences; RMIT University
- Dr Sabira Shrestha, National Centre for Immunisation Research and Surveillance (NCIRS)
- Vicky Jacobson, Coordinator, Refugee Health Network Queensland
- Dr Carissa Bonner, The School of Public Health, University of Sydney

#### **Special thanks:**

We would like to thank the following people for reviewing the glossary

- Associate Professor Christopher Blyth, Wesfarmers Centre of Vaccines and Infectious Diseases, Telethon Kids Institute.
- Dr Nadia Chaves
- Benine Muriithi, Patricia Argüello de Avila and Mariam Elliott, Refugee Health Network Queensland
- Fartih Karakas and Ismail Akinci, All Graduates
- Dr Jan Fizzell, Senior Medical Advisor, Public Health Response Branch, NSW Health

#### Thanks to

The Health Literacy Lab for testing the glossary using their online real-time editor (*Ayre, J., et al. (2021). Sydney Health Literacy Lab (SHLL) Health Literacy Editor*). Available at https://hdl.handle.net/2123/24642

## Contents

<u>A</u>	4	Μ	16
В	6	N	17
C	7	0	18
D	9	Ρ	19
E	10	R	21
G	11	S	22
H	12	<u>T</u>	23
<u>I</u>	13	V	24
L	15	W	26

Α	DEFINITION	SIMPLIFIED DEFINITION
Adjuvant	A vaccine component distinct from the antigen that enhances the immune response to the antigen.	An adjuvant is an ingredient used in some vaccines. It helps our bodies make a stronger immune response. The adjuvant works together with other parts of the vaccine. They have been used in some vaccines for over 70 years.
Adverse event (reaction)	An unexpected, unwanted, or serious reaction to a medication or vaccine.	Any unexpected or serious effect that happens after a vaccine or medicine. Something that was not expected to happen.
Adverse event following immunisation (AEFIs)	Any untoward medical occurrence that follows immunisation. It does not necessarily have a causal relationship with the vaccine	An unexpected effect that happens after vaccination. The vaccine may have not been the reason for the problem.
Advisory Committee on Vaccines (ACV)	The ACV provides independent medical and scientific advice to the Australian Government's Minister for Health and the Therapeutic Goods Administration (TGA) on issues relating to the safety, quality and efficacy of vaccines supplied in Australia including issues relating to pre-market assessment, post-market monitoring and safe use in national immunisation programs.	A group of experts that gives medical and scientific advice. The group talks to the Australian Government's Minister for Health and the Therapeutic Goods Administration (TGA). They give advice on issues about vaccine safety and use.
Allergy	An immune response to a foreign substance that is harmless in most people. The foreign substance could be a food, medicine or vaccine component.	An immune response to a foreign material that is harmless to most people. The foreign material could be food, medicine or a part of a vaccine.
Anaphylaxis	An immediate and severe allergic reaction to a substance (e.g. food or drugs). Symptoms of anaphylaxis include breathing difficulties, loss of consciousness and a drop in blood pressure. This condition can be fatal and requires immediate medical attention.	A quick and serious allergic reaction. This could be a reaction to food or medicine. Symptoms can include breathing difficulties, loss of consciousness and a drop in blood pressure. The person will need urgent medical attention and can sometimes die.
Antibody	A protein found in the blood that is produced in response to foreign substances (e.g., viruses) invading the body. Antibodies protect the body from disease by binding to these organisms and destroying them.	When the body gets sick or gets a vaccine, the body will make antibodies to protect it against that disease. The body can then recognise the germs when that same disease happens again.



A Cont'd	DEFINITION	SIMPLIFIED DEFINITION
Antigen	A foreign substance which is detected by the immune system. The presence of antigens in the body triggers an immune response, usually the production of antibodies.	A foreign (external) substance like bacteria, viruses, or fungi that cause infection and disease if they get inside the body . The immunes system detects them and produces antibodies to fight them.
Association	The degree to which the occurrence of two variables or events is linked. Association describes a situation where the likelihood of one event occurring depends on the presence of another event or variable. However, an association between two variables does not necessarily imply a cause- and-effect relationship. The term association and relationship are often used interchangeably	A link between one event taking place at the same time as another event. The fact that they are happening together does not prove that one event caused the other event.
Asymptomatic	A person that is not showing symptoms	Someone with no sign of infection.
Attenuated vaccine	An attenuated vaccine (or a live attenuated vaccine) is a vaccine created by reducing the virulence of a pathogen but keeping it viable (or "live"). Attenuation takes an infectious agent and alters it so that it becomes harmless or less dangerous. These vaccines contrast to those produced by "killing" the virus (inactivated vaccine).	Live vaccines use a less strong (or attenuated) form of the germ that causes a disease. These vaccines are like the natural infection that they help prevent. They create a strong and long-lasting immune response.
Australian Immunisation Register	It is a national register that records all vaccines given to all people in Australia	An electronic register that contains information on all vaccines given to all Australians.
Australian Technical Advisory Group on Immunisation (ATAGI)	ATAGI's role is to advise the Minister for Health on the medical administration of vaccines available in Australia.	A group of experts that helps the Government to make decisions on the use of vaccines in Australia.



В	DEFINITION	SIMPLIFIED DEFINITION
Batch assessment	The Therapeutic Goods Administration (TGA) ensures there is an independent quality assessment of every batch of vaccine supplied in Australia.	A process of checking that the vaccines used in Australia are of high quality. The Therapeutic Goods Administration will do these checks.
Boost (Booster injection)	An additional dose of a vaccine that re-stimulates the immune after the effects of an earlier dose wear off. It is not yet clear whether booster shots of a COVID-19 vaccine will be necessary.	An additional vaccine after the first one, given to either build up better immunity or to make sure the immunity lasts longer.
Breakthrough infection	Development of a disease despite a person's having responded to a vaccine.	When someone can get the infection even though they are vaccinated.

С	DEFINITION	SIMPLIFIED DEFINITION
Cell culture	A laboratory technique that can be used to make some vaccines, where cells can be grown in a nutrient-rich liquid to either I) make virus to be inactivated for inactivated vaccines, ii) make antigen for protein subunit vaccines or iii) make viral vectors.	Using cells grown in liquid to make vaccine ingredients.
<b>Clinical Trial</b>	A research study in which one or more human subjects are progressively assigned to one or more interventions (which may include placebo/sham or other control) to evaluate the effects of those interventions on health-related biomedical or behavioural outcomes.	A type of research study. People either receive a new vaccine or are in the control group. The control group may receive a different vaccine or a placebo, meaning a simple substance with no effects on the body. Participants usually do not know which group they are in. Scientists test the safety and benefits of new vaccines.
Co-administration	Co-administration of vaccines is defined as the administration of more than one vaccine on the same day.	To be given more than one vaccine on the same day.
Coalition for Epidemic Preparedness Innovations (CEPI)	One of the international organisations that established the COVID-19 Vaccine Global Access (COVAX) Facility. The facility enables participating governments, including lower income countries, to access safe and effective vaccines for 20 per cent of their population.	An international organisation that will help many countries gain access to COVID-19 vaccines. It will help governments, including lower income countries, to access safe and effective vaccines for 20% of their population.
Cold chain	The system of transporting and storing vaccines within the safe temperature range. This is normally between +2°C to +8°C.	Shipping and storing vaccines at the correct temperature.
Combination vaccine	A product containing components that can be divided equally into independently available routine vaccines.	Combination vaccines take two or more vaccines that could be given individually and put them into one shot.
Conjugate vaccine	A vaccine made by chemically linking a protein molecule with a tiny sugar compound. The sugar compound comes from the cell coating of the bacteria. This improves the person's immune response to the vaccine. Example: Meningococcal vaccine	The joining together of two compounds (usually a protein and polysaccharide) to increase a vaccine's effectiveness.



C Cont'd	DEFINITION	SIMPLIFIED DEFINITION
Contraindication	A condition in a person wanting to be vaccinated that increases the risk for a serious adverse reaction.	An illness (or health condition) that increases the risk for a serious adverse health consequence.
Control group	A control group is an untreated research sample against which all other groups or samples in the research is compared. A control group is constructed by randomly assigning people to either the control group or to one or more "treatment" groups.	A group of people who do not receive the vaccine or drug being tested. Instead, they may get the normal intervention (drug, vaccine, or treatment), a placebo or nothing. The aim of the trial is to compare what happens in each group. The results must be different enough between the two groups to prove that the difference has not just occurred by chance. A placebo is a 'dummy' treatment, such as a sugar pill, that looks the same.
Convalescent plasma	Blood plasma that is obtained from an individual who has recovered from an infectious disease and contains antibodies against the infectious agent of the disease and may be administered by intravenous transfusion to prevent or treat infection in other individuals.	Plasma is the liquid part of blood. It is collected from a person after they have had an infection. The liquid contains antibodies against the germ. Sometimes this plasma can be given to other people to prevent them getting sick or to help them get better.
COVAX	COVAX is co-led by Gavi, the Coalition for Epidemic Preparedness Innovations (CEPI) and World Health Organisation (WHO). Its aim is to accelerate the development and manufacture of COVID-19 vaccines, and to guarantee fair and equitable access for every country in the world.	An international partnership that aims to support the development and delivery of the COVID-19 vaccines fairly around the world.



D	DEFINITION	SIMPLIFIED DEFINITION
Deltoid	A muscle in the upper arm where vaccines are usually administered given.	A muscle in the upper arm where vaccine is given
Dose	A quantity of a medicine or drug taken or recommended to be taken at a particular time.	An amount of a medicine or drug taken
Dose/vaccination schedule	This is a timetable of vaccines that an individual (baby, infant, adolescent adult, elderly) is to receive throughout their lifetime.	A list of times to have vaccines throughout your life.
Dosing error	When medications are administered in the wrong amounts, at the wrong frequency or to the wrong person	When medicines are given in the wrong amount, at the wrong time point or to the wrong person



G	DEFINITION	SIMPLIFIED DEFINITION
Guillain-Barré syndrome	An autoimmune condition that causes muscle weakness, numbness or paralysis.	A rare event where a person's immune system attacks the nerves and can cause muscle weakness, numbness or not being able to move part or most of the body.

Η	DEFINITION	SIMPLIFIED DEFINITION
Herd immunity	This is a type of infection control that occurs naturally, or through immunisation programs, when a large enough portion of a population becomes immune to a disease to stop further spread. Immunity may be either by recovering from the disease or by being vaccinated against it. In the case of COVID-19, the possibility of herd immunity remains unclear due to the uncertainty of long term immunity to this virus.	When most people in a community have protection against an infection. High levels of protection make it more and more difficult for the germ to pass from person to person. This can successfully stop the spread of disease in the community.



	-	~	
(			1
(		٦	J
1	_	~	

I Cont'd	DEFINITION	SIMPLIFIED DEFINITION
Immunocompromised	Someone whose immune system struggles to fight off diseases or infections as well as most people. There are several illnesses which can cause someone to become immunocompromised. Immunocompromised people are at a higher risk from COVID-19.	Someone whose immune system may not be able to fight off an illness as well as other people. There are several illnesses that can cause someone to become immunocompromised.
Immunogenicity	The ability of a particular substance, such as a vaccine, to provoke an immune response, or the degree of response it provokes.	The power of a substance to produce an immune response. The substance can be present in a vaccine.
Inactivated vaccine	A vaccine made from viruses and bacteria that have been killed through physical or chemical processes. These killed organisms cannot cause disease.	A vaccine made from a germ that has been killed. The germ is killed either by high heat or by chemicals. When this killed germ is injected into your body, it helps your immune system learn to find the germ, without the risk of getting sick.
Incidence	This term is used to describe the rate of new illness or injuries (morbidities) within a precise population over a specified period of time. Incidence can mean the number of new cases in a community, or the number of new cases in a unit of population over a period of time.	Incidence can mean the number of new disease cases in a town, city, state or country or the number of new cases in a particular group of people over a certain amount of time.
Infection	When bacteria or viruses invade the body. If the body cannot fight the infection, it may cause disease.	When a germ (bacteria or virus) enters the body. If the body cannot fight the growth of germs, it may cause disease.
Injection site reaction	Injection site reactions are the most common adverse events following immunisation. These include pain, itching, swelling or redness around the site of injection. These reactions are usually mild and last for 1–2 days.	The most common side-effects after a vaccine. This can include pain, itching, swelling or redness around the site of injection. These are usually mild and last for 1–2 days.
Intramuscular injection	An injection into the muscle. Vaccines are usually injected into a muscle in the upper outer thigh or upper arm.	Where the vaccine is given into the muscle.



L	DEFINITION	SIMPLIFIED DEFINITION
Lipid	Essentially a type of fat. Lipids are being used to make a protective bubble around mRNA in mRNA vaccines to prevent it from being broken down before it enters a cell.	Lipid is fat that is used to make a protective bubble around the mRNA in mRNA vaccines. mRNA is a type of small molecule. mRNA is very weak and breaks down quickly in the body if it is not protected. Once the mRNA is transported into the cell, it is broken down inside the cell.
Live vaccine	A vaccine containing live viruses or bacteria that are weakened (attenuated) to produce an immune response in the recipient without causing the serious effects of the disease.	The vaccine uses a live form of the bacterium/virus that may be weakened.
Long-COVID (also known as post-covid condition)	'Long COVID' is a condition where people with COVID-19 experience persistent symptoms that continue for at least 2 months. Common symptoms include fatigue, shortness of breath, cognitive dysfunction ('brain fog') and others. These symptoms generally affect the person's everyday functioning. Symptoms can fluctuate or relapse over time.	When people with COVID-19 have symptoms for two months or more. Common symptoms include feeling tired, shortness of breath, memory problems, and others. These symptoms can affect the person's daily functioning.

Μ	DEFINITION	SIMPLIFIED DEFINITION
Messenger RNA (mRNA)	An RNA produced by transcription that carries the code for a particular protein from the nuclear DNA to a ribosome in the cytoplasm and acts as a template for the formation of that protein.	A type of small molecule that your cells use as instructions to make protein. mRNA tells your cells how to put together a specific protein using building blocks (called amino acids). You have many millions of mRNA molecules in your body at any one time- all being used to make proteins.
Minimum interval	The shortest amount of time required between 2 doses of a vaccine to provide an adequate immune response to the 2nd dose. If the interval between the doses is shorter than the minimum interval, the vaccine may not be effective.	The amount of time a person needs to leave between receiving their 1st and 2nd vaccines.
Mixed (heterologous) schedules	In some circumstances, an alternative formulation, brand or vaccine platform may be recommended for the second dose (if not contraindicated).	Sometimes a different mixture, brand or vaccine platform may be given for the second vaccine.
Morbidity	Morbidity is the state of having a specific illness or condition.	Illness that happens due to a specific infection or condition.
Mortality	The number of deaths that have occurred due to a specific illness or condition.	Deaths that happen due to a specific infection or condition.
mRNA vaccine	Contain material from the virus that causes COVID-19 that gives our cells instructions for how to make a harmless protein that is unique to the virus. After our cells make copies of the protein, they destroy the genetic material from the vaccine. Our bodies recognize that the protein should not be there and build T-lymphocytes and B-lymphocytes that will remember how to fight the virus that causes COVID-19 if we are infected in the future.	mRNA vaccines teach our cells how to make a harmless protein—or even just a piece of a protein. This protein activates an immune response inside our bodies. That immune response, which produces antibodies, is what protects us from getting very unwell if the real virus enters our bodies.
Multi-dose vial	Multi-dose vials contain more than one dose of a medicine/vaccine in a single vial. Vial: a small container, typically cylindrical and made of glass, used especially for holding liquid medicines.	The containers (vials) hold more than one dose of a medicine or vaccine in a single vial.
Myocarditis	An inflammatory disease of the heart muscle.	A disease where the heart muscle becomes inflamed.

Ν	DEFINITION	SIMPLIFIED DEFINITION
National Immunisation Programme (NIP)	A programme in Australia that funds free vaccines for eligible people to increase immunization rates and help reduce vaccine-preventable diseases.	The vaccines that are given for free in Australia to children, adolescents and adults.
Neutralisation	One way that our immune system can protect us from infection is neutralisation, where our immune cells make a protein called an antibody that coats the virus and prevents it from getting into our cells. They also help other parts of the immune system recognise and destroy the virus.	One way that our immune system can protect us from an infection. Our immune system makes antibodies that stick all over the surface of a virus. When the virus tries to stick onto our cells, the antibodies get in the way and stop the virus from getting into our cells. They also help other parts of the immune system recognise and destroy the virus

0	DEFINITION	SIMPLIFIED DEFINITION
Outbreak	The occurrence of cases of disease in excess of what would normally be expected in a defined community, geographical area, and/or season, but in lower than epidemic numbers.	When the number of disease cases is higher than what has previously happened in a city or country, or in a group of people or time of year.

Ρ	DEFINITION	SIMPLIFIED DEFINITION
Pandemic	Worldwide spread of a new disease, such as a new influenza virus or the coronavirus, COVID-19.	Spread of a new disease to every country around the world.
Paediatric inflammatory multisystem syndrome temporally associated with SARS-CoV-2 (PIMS-TS)	Paediatric multisystem inflammatory syndrome – temporally associated with SARS-CoV-2 (or PIMS-TS) is the current name given to the rare hyperinflammatory state seen in children who have been infected with SARS-CoV-2.	This is the name of a rare illness in some children after having COVID-19 where parts of the body becomes larger than normal.
Paraesthesia	Paraesthesia is an unusual feeling in the skin, such as tingling or a crawling sensation.	A feeling in the skin that is not normal, like feeling something crawl under the skin or tingling or numbness.
Passive immunisation	The transfer of antibodies from one person to another, providing temporary protection from diseases. Disappears within a few weeks or a month.	Occurs when a person is given antibodies to a disease rather than producing them through his or her own immune system. Such as a newborn baby receiving antibodies from their mum.
Pathogen	An agent of disease such as a virus or bacterium.	A germ that can cause disease if you are infected, such as a virus
Peer-review	A process where independent scientists examine findings from a study and determine if the work has been performed well and the findings are supported by the data.	Independent experts examine other people's research to make sure it is appropriate and correct.
Pericarditis	An inflammatory disease of the lining of the heart muscle.	A disease where the lining of the heart muscle becomes inflamed.
Placebo	A substance or treatment that has no effect on human beings.	A substance or treatment that has no effect on human beings.
Polysaccharide vaccine	Vaccines that are composed of long chains of sugar molecules that resemble the surface of certain types of bacteria. Polysaccharide vaccines are available for pneumococcal disease.	A vaccine containing long threads of sugar molecules, which look like the surface of some kinds of bacteria. Polysaccharide vaccines are available for pneumococcal disease (such as pneumonia).

P Cont'd	DEFINITION	SIMPLIFIED DEFINITION
Post-market vaccine effectiveness studies (also known as 'real world' studies or Phase IV clinical trials)	These studies are designed to monitor the effectiveness of the approved intervention in the general population and to collect information about any adverse effects associated with widespread use over longer periods of time.	These types of studies are used to see whether the approved vaccine works in the real world and to look for any side effects.
Pre-Clinical Trial	A research study that is done prior to a Clinical Trial using cells or using animals to test whether a vaccine is promising enough to be evaluated with human volunteers.	A research study done before a clinical trial. The study tests whether a vaccine is safe to test on humans. As part of the COVID-19 trials, animal models included experiments on animals including mice and macaques.
Prevalence	The proportion of a population that have a particular disease in a certain period of time.	The total number of people in a population that have a disease at a certain time.
Primary course	A primary course of a vaccine is the number of doses it takes to achieve a good level of protection against disease.	The number of vaccines it takes to achieve a good level of protection against disease.
Prime	The first time a vaccine is given	The first time a vaccine is given
Protein subunit vaccine	Vaccines that include harmless pieces of a virus instead of the entire germ. Once vaccinated, our immune system recognizes that the proteins do not belong in the body and begins making T-lymphocytes and antibodies. If we are ever infected in the future, memory cells will recognize and fight the virus.	Include harmless pieces (proteins) of the germ instead of the entire germ. Once vaccinated, our bodies recognize that the protein should not be there and build blood elements called T-lymphocytes and antibodies that will remember how to fight the germ if we are exposed in the future.

/	7		1
(1	٢		t)
1		_	/

R	DEFINITION	SIMPLIFIED DEFINITION
Reactogenicity	The physical manifestation of the inflammatory response that develops to vaccination, and can include injection-site pain, redness, swelling or induration at the injection site, as well as systemic symptoms, such as fever, myalgia, or headache.	A group of effects that often happen after vaccination. It can include pain, redness or swelling around where the vaccine was injected. A person might feel tired, or hot or have a headache. Importantly, these are signs that an immune response is working.
Regulatory body	A government organisation that decides which vaccines are able to be registered in a country and legally supplied to people in the country.	A government organisation that decides which vaccines can be registered in a country and legally used in the country.
Roll out	Officially launch or introduce a new product or service. Australia's COVID-19 vaccine national roll-out strategy includes the following phases: 1a, 1b, 2a, 2b, 3.	The introduction of a new drug or vaccine. For the COVID-19 vaccination program this includes multiple phases: 1a, 1b, 2a, 2b, 3. Priority groups are identified by considering current public health and medical evidence on who would be most affected if they got COVID-19.

S	DEFINITION	SIMPLIFIED DEFINITION
SARS-CoV-2	The official name of the virus that causes the disease known as COVID-19. It is part of a bigger family of viruses called coronaviruses.	The official name of the virus that causes the disease known as COVID-19. It belongs to family of viruses called coronaviruses.
Serology	Measurement of antibodies, and other immunological properties, in the blood serum.	Measuring the level of antibodies (immune proteins) present in the blood.
Side Effect	Undesirable reaction resulting from immunization.	Any unwanted or unexpected effects of a vaccine.
Spike protein	A glycoprotein that protrudes from the envelope of some viruses (such as a coronavirus) and facilitates entry of the virion into a host cell by binding to a receptor on the surface of a host cell followed by fusion of the viral and host cell membranes.	Coronaviruses have sharp bumps on their surface. Those bumps are called spike proteins. They help the virus enter a person's cells.



Т	DEFINITION	SIMPLIFIED DEFINITION
Therapeutic Goods Administration (TGA)	The TGA is responsible for assessing vaccines and other medicines before they can be used in Australia	The Therapeutic Goods Administration (TGA) is responsible for checking vaccines and other medicines before they can be used in Australia.
Thrombosis with thrombocytopenia syndrome (TTS)	A rare and new syndrome. The condition involves blood clots occurring in body sites like the brain or abdomen) together with low platelet levels.	A newly described serious condition. A person gets unusual blood clots in the brain or in other parts of the body. It is also associated with low platelet levels.
Transmission	The ability of a virus to pass from one person to another.	The ability of a virus to pass from one person to another.

V	DEFINITION	SIMPLIFIED DEFINITION
Vaccination	The process of administering a vaccine to the body. This is typically done through an injection	Giving a vaccine to help the immune system develop protection from a specific disease. Commonly used terms include shot, jab, needle, and inoculation.
Vaccine	Medicines that help prepare our immune systems to defend against infection from certain diseases. Usually, vaccines are given before the person is exposed to the disease. Each vaccine stimulates the immune system to make antibodies against a particular virus or bacteria. Some vaccines provide lifelong immunity, but others may require 'booster shots' to maintain immunity.	A type of medicine that supports our immune system to fight against certain germs and prevent disease. Usually, vaccines are given before the person encounters the germ. Each vaccine promotes the immune system to make antibodies against the germ.
Vaccine Candidate	An experimental vaccine that is still being tested.	A new vaccine that is still being tested and is not licensed.
Vaccine hesitancy	Refers to delay in acceptance or refusal of vaccines despite availability of vaccine services.	When a person is unsure about a vaccine and delays or refuses to receive the available vaccine.
Variant (mutation)	Tiny changes in the virus that can occur to the genetic information that occur during the process of replication	Tiny changes in the genetic information inside a virus. Variants can occur when a virus multiplies or makes copies of itself.
Vial	A small container used to hold medicine	A small container used to hold medicine.
Viral shedding	Viral shedding occurs when a virus replicates inside your body and is released into the environment. At that point, it may be contagious.	When the virus made inside your body starts to be released into your surroundings. At that point, it may be spread or passed on to other people.

V Cont'd	DEFINITION	SIMPLIFIED DEFINITION
Viral vector vaccine	Contain a weakened version of a live virus — a different virus than the one that causes COVID-19 — that has genetic material from the virus that causes COVID-19 inserted in it (this is called a viral vector). Once the viral vector is inside our cells, the genetic material gives cells instructions to make a protein that is unique to the virus that causes COVID-19. Using these instructions, our cells make copies of the protein. This prompts our bodies to build T-lymphocytes and B-lymphocytes that will remember how to fight that virus if we are infected in the future.	Contains a modified version of a different virus from the one that causes COVID-19. Inside the modified virus, there is material from the virus that causes COVID-19. This is called a "viral vector." Once the viral vector is inside our cells, the genetic material gives cells instructions to make a protein that is unique to the virus that causes COVID-19. Using these instructions, our cells make copies of the protein. This prompts our bodies to build T-lymphocytes and B-lymphocytes that will remember how to fight that virus if we are infected in the future.



W	DEFINITION	SIMPLIFIED DEFINITION
Waning immunity	The loss of protective antibodies over time.	When your level of immunity gets lower and lower with time.